

Deformed Distance Duality Relations and Supernovae Dimming

J. A. S. Lima^{1,a}, J. V. Cunha^{2,b}, V. T. Zanchin^{3,b}

^a*Departamento de Astronomia, Universidade de São Paulo, USP,
05508-900 São Paulo, SP, Brazil*

^b*Centro de Ciências Naturais e Humanas, Universidade Federal do ABC
Rua Santa Adélia 166, 09210 - 170, Santo André, SP, Brazil*

ABSTRACT

The basic cosmological distances are linked by the Etherington cosmic distance duality relation, $\eta(z) = D_L(z)(1+z)^{-2}/D_A(z) \equiv 1$, where D_L and D_A are, respectively, the luminosity and angular diameter distances. In order to test its validity, some authors have proposed phenomenological expressions for $\eta(z)$ thereby deforming the original Etherington's relation and comparing the resulting expressions with the available and future cosmological data. The relevance of such studies is unquestionable since any violation of the cosmic distance duality relation could be the signal of new physics or non-negligible astrophysical effects in the usually assumed perfectly transparent Universe.

In this letter, we show that under certain conditions such expressions can be derived from a more fundamental approach with the parameters appearing in the $\eta(z)$ expression defining the cosmic absorption parameter as recently discussed by Chen and Kantowski. Explicit examples involving four different parametrizations of the deformation function are given. Based on such an approach, it is also found that the latest Supernova data can also be explained in the framework of a pure cold dark matter model (Einstein-de Sitter). Two different scenarios with cosmic absorption are discussed. Only if the cosmic opacity is fully negligible, the description of an accelerating Universe powered by dark energy or some alternative gravity theory must be invoked.

Subject headings: Supernovae:general, cosmic distances, duality relation

¹limajas@astro.iag.usp.br

²cunhajv@astro.iag.usp.br

³zanchin@ufabc.edu.br

1. Introduction

The *cosmic distance duality relation* (CDDR) is a mathematical identity relating the *luminosity distance* D_L with the *angular diameter distance* D_A by the expression:

$$\frac{D_L}{D_A}(1+z)^{-2} = 1. \quad (1)$$

The validity of this constraint uniting the two basic distances in cosmology depends neither on the Einstein field equations nor on the nature of the matter-energy content. It only requires the phase space conservation of photons and that sources and observers are connected by null geodesics in a Riemannian spacetime. Therefore, it remains valid for spatially homogeneous and isotropic (anisotropic) cosmologies, as well as for inhomogeneous cosmological models (Etherington 1933, Basset and Kunz 2004).

The above relation is usually taken for granted when relativistic models of the Universe are confronted to the existing cosmological observations. Despite that, the distance-duality relation is in principle testable by means of astronomical observations. The basic idea is to find cosmological sources whose intrinsic luminosities are known (standard candles) as well as their intrinsic sizes (standard rulers). After determining both D_L and D_A at the same redshift it should be possible to test directly the Etherington result. Naturally, by cosmological sources with known D_L and D_A we are not referring necessarily to the same class of objects. Under certain conditions, as recently discussed by Holanda et al. (2010), one may consider two different classes of objects as, for instance, supernovae and galaxy clusters for which D_L and D_A are separately determined. Note also that ideally both quantities must be measured in such a way that any relationship coming from specific cosmological models are not used, that is, they must be determined by means of intrinsic astrophysically measured quantities. In practice, the validity of the CDDR has been tested by assuming a phenomenological deformed expression of the form (Holanda et al. 2010, Meng et al. 2011, Nair et al. 2011, Khedekar & Chakabarti 2011, Gonçalves et al. 2011):

$$\frac{D_L}{D_A}(1+z)^{-2} = \eta(z), \quad (2)$$

where $\eta(z)$ is the deformation function which quantifies a possible epoch-dependent departure from the standard photon conserving scenario ($\eta = 1$).

As it appears, a deformed CDDR can also be adopted to test the possibility of a new physics. In this line, Basset & Kunz (2004) used supernovae type Ia (SNe Ia) data as measurements of the luminosity distance and the estimated D_A from FRIB radio galaxies (Daly & Djorgovski 2003) and ultra compact radio sources (Gurvitz 1999; Lima & Alcaniz

2000, 2002; Santos & Lima 2008) in order to test possible new physics signatures based on the following expression:

$$I. \quad \eta(z) = (1+z)^{\beta-1} \exp \left[\gamma \int_0^z \frac{dz'}{E(z')(1+z')^\delta} \right], \quad (3)$$

where $E(z') \equiv H(z')/H_0$ is the dimensionless Hubble parameter (H_0 is the Hubble constant). Note that for arbitrary values of δ , the strict validity of the DD relation corresponds to $(\beta, \gamma) \equiv (1, 0)$. By marginalizing on Ω_M , Ω_Λ and Hubble parameters, they found a 2σ violation caused by excess brightening of SNIa at $z > 0.5$. It was also argued that such an effect would be associated to lensing magnification bias.

De Bernardis, Giusarma & Melchiorri (2006) also searched for deviations of the standard CDDR by using the angular diameter distances from galaxy clusters provided by the sample of Bonamente *et al.* (2006). By assuming $\eta(z) = \text{constant}$, they obtained a non violation of CDDR in the framework of the cosmic concordance Λ CDM model. Later on, Avgoustidis et al. (2009, 2010) working in the context of a flat Λ CDM model also adopted an extended CDDR expressed as

$$II. \quad \eta(z) = (1+z)^\epsilon. \quad (4)$$

The above deformation function is a particular case ($\beta = 1 + \epsilon, \gamma = 0$) of the general expression adopted by Basset and Kunz (2004). In their analysis, the recent SNe Ia data as compiled by Kowalski et al. (2008) were combined with the latest measurements of the Hubble expansion at redshifts in the range $0 < z < 2$ (Stern et al. 2010), and the free parameter was constrained to be $\epsilon = -0.04^{+0.08}_{-0.07}$ (2σ). More recently, such a parametrization has also been adopted by Khedekar & Chakabarti (2011) in their studies connecting the Tolman test and CDDR using the redshifted 21 cm wavelength from disk galaxies. It was also argued that future data from the planned Square Kilometer Array (SKA) may provide the best test to detect any violation of the cosmic distance duality relation.

In a series of papers, Holanda, Lima & Ribeiro (2010, 2011, 2011a) also explored a different route to test the CDDR based on the following deformation functions:

$$III. \quad \eta(z) = 1 + \eta_0 z, \quad IV. \quad \eta(z) = 1 + \eta_0 z / (1 + z). \quad (5)$$

A basic difference between the first and the second parameterization is that the later includes a possible epoch dependent correction which avoids the divergence at extremely high z . At low redshifts, when second order terms are neglected, the second parametrization reduces to the first one, that is, $\eta(z) \simeq 1 + \eta_0 z$. Such one-parametric formulas are also interesting because in the limit of extremely low redshifts ($z \ll 1$), one finds $\eta(z) = 1$ as should be expected since $D_L = D_A$ at this limit (see also parametrization II). In addition, for a given

data set, the likelihood of η_0 must be peaked at $\eta_0 = 0$, in order to satisfy the standard duality relation. In this context, by taking the SNe Ia from Constitution data (Hicken et al. 2009) and galaxy clusters samples compiled by De Fillipis et al. (2005) and Bonamente et al. (2006), a direct test of the CDDR was accomplished (Holanda et al. 2010). As an extra bonus, the consistency between the strict validity of CDDR and the assumptions about the geometry based on elliptical and spherical β models was detailedly discussed. The sphericity assumption for the cluster geometry resulted in a larger incompatibility with the validity of the duality relation in comparison with an isothermal non spherical cluster geometry. More recently, such expressions were adopted by Gonçalves et al. (2011) in their studies on the validity of CDDR by including data from X-ray gas mass fraction of galaxy clusters.

In this connection, it is also worth mentioning that Li, Wu & Yu (2011) rediscussed this independent cosmological test by using the latest Union2 SNe Ia data and the angular diameter distances from galaxy clusters thereby obtaining a more serious violation of the standard duality expression. In a simultaneous but independent work, Nair, Jhingan & Jain (2011) also investigated the strict validity of the CDDR by using the latest Union2 SNe Ia data and the angular diameter distances from galaxy clusters, FRIIb radio galaxies and mock data. As an attempt to determine a possible redshift variation of the CDDR relation, they proposed six different (one and two indexes) parametrizations including, as particular cases, the ones adopted by Holanda et al. (2010, 2011a). As physically expected, their results depend both on the specific parametrization and the considered data sample. In particular, they conclude that the one index parametrizations, namely: $\eta_V = \eta_8/(1+z)$ and $\eta_V = \eta_9 \exp\{[z/(1+z)]/(1+z)\}$, does not support the CDDR relation for the given data set. Meng, Zhang & Zhan (2011) also reinvestigated the model independent test by comparing two different methods and several parametrizations (one and two indexes) for $\eta(z)$. Their basic conclusion is that the triaxial ellipsoidal model is suggested by the model independent test at 1σ while the spherical β model can only be accommodated at 3σ confidence level thereby agreeing with the results earlier derived by Holanda et al. (2010).

It should be stressed that all the above described attempts to test the CDDR have been carried out based on a phenomenological approach. Usually, it is also not clear whether the nonstandard relation is the result of a modified luminosity distance or whether it should be associated to an extended angular diameter distance or both (see, however, Basset and Kunz 2004). At this point, one may also ask whether such expressions to $\eta(z)$ can be derived from a more fundamental approach. In the affirmative case, it is also important to study their consequences for the present accelerating stage of the Universe.

In this letter we consider both questions. Firstly, we show how any deformed CDDR can be derived by analysing possible theoretical modifications on the luminosity distance without

refraction effects. Analytical expressions for the dimensionless cosmic absorption parameter describing the above four parametrizations will be explicitly obtained in the framework of Gordon’s optical metric as developed by Chen and Kantowski (2009a, 2009b) to include cosmic absorption. It will be also explicitly assumed that the angular diameter distances are not modified because their measurements involve only standard rulers and angular scales, and, more important, possible refractive effects have been neglected. Apart such hypotheses, the approach discussed here is quite general and can be applied for any deformation function, $\eta(z)$. Secondly, we also apply our results for the latest SNe Ia data. As we shall see, the modified luminosity distance can accomodate the observed supernova dimming even for a non-relativistic cold dark matter (CDM) Einstein - de Sitter model ($\Omega_M = 1$). The validity of the Λ CDM description is obtained only in the extreme limit of perfect cosmic transparency (negligible cosmic absorption).

2. Luminosity Distance and Duality Relation

The concept of an optical metric was introduced long ago in a seminal paper by Gordon (1923). He proved the existence of a mapping between any solution of the general relativistic Maxwell’s equations for a fluid with refraction index $n(x)$ and the vacuum solutions in a related optical spacetime. In Gordon’s treatment, only the refraction phenomenon was considered. More recently, by describing the Maxwell field as a monochromatic wave, Chen and Kantowski (2009a, 2009b) generalized such treatment in order to include the possibility of an absorption phenomenon in the Universe. As it appears, the presence of such an effect breaks naturally the validity of the standard distance duality relation as given by Eq. (1) since the light absorption violates the photon number conservation law. In this case, they prove that the luminosity distance takes the following form:

$$D_L(z) = e^{\tau/2} D_L^S(z) = \frac{e^{\tau/2}(1+z)}{H_0} \int_0^z \frac{dz'}{E(z')}, \quad (6)$$

where τ denotes the optical depth associated to the cosmic absorption of the Universe and the superscript S specify the standard luminosity distance (no absorption) for which the Universe is assumed to be transparent. It is worth noticing that effects coming from a possible new physics like the interaction between photons and dark matter as discussed by Basset and Kunz (2004) are assumed to be negligible (no new physics takes place). For a spatially homogeneous and nondispersive (grey) absorption, the quantity τ as derived by Chen & Kantowski (2009a) reads

$$\tau(z) = \int_0^z \frac{\alpha_* dz'}{(1+z')E(z')}, \quad (7)$$

where α_* is the dimensionless cosmic absorption parameter (α/H_0 in the notation of Chen & Kantowski). In the above expression the refractive index was fixed to unity ($n(z)=1$), and, therefore, any possible refraction effect has been neglected. This is an important point, since in this case the standard angular diameter distances are not modified.

Let us now discuss how the several functions $\eta(z)$ defining the deformed CDDR introduced in an ad-hoc way can be related with the dimensionless cosmic absorption parameter, $\alpha_*(z)$. As one may check, by inserting expression (6) into (2), we obtain that the cosmic optical depth and the deformation function must be related by the simple expression $e^{\tau/2} = \eta(z)$. In addition, this also means that any smooth $\eta(z)$ deformation function defines an effective cosmic absorption parameter given by:

$$\alpha_*(z) = \frac{2\eta'(z)(1+z)E(z)}{\eta(z)}, \quad (8)$$

where a prime denotes derivative with respect to the reshift. Such a relation uniting the dimensionless cosmic absorption parameter and the deformation function, usually introduced by hand in the distance duality relation, is one of the main results of this section. When $\eta(z)$ is constant the cosmic absorption $\alpha_*(z)$ is identically null, and, therefore, $\tau = 0$. As remarked earlier, this also means that only the standard relation as determined by Etherington (1933) is possible at this limit, that is, $\eta(z) = 1$ (note that τ and η are related by an exponential function). As one may check, the four deformed distance duality relations can analytically be expressed in terms of the dimensionless absorption parameter as:

$$\text{I. Basset and Kunz (2004): } \alpha_*(z) = 2[(\beta - 1)E(z) + \gamma(1+z)^{1-\delta}],$$

$$\text{II. Avgoustidis et al. (2009): } \alpha_*(z) = 2\epsilon E(z),$$

$$\text{III. Holanda et al. (2010): } \alpha_*(z) = \frac{2\eta_0(1+z)E(z)}{1+\eta_0 z},$$

$$\text{IV. Holanda et al. (2010): } \alpha_*(z) = \frac{2\eta_0 E(z)}{1+z+\eta_0 z}.$$

We see that the dimensionless Hubble parameter, $E(z)$, and the free parameters appearing in the deformation functions define completely the cosmic absorption parameter as introduced by Chen and Kantowski (2009a, 2009b).

At this point some comments are in order. Initially, we notice that in the limiting case $\gamma = 0$ and $\beta = \epsilon + 1$, the first expression for the cosmic absorption reduces to the second one. This should be expected since at such a limit the general deformation function of Basset and Kunz (2004) reduces to the one proposed by Avgoustidis (2009). Note also that such

expressions are satisfied for all Friedmann-Robertson-Walker (FRW) geometries (arbitrary values of Ω_k) and energetic content (baryons, dark matter, and dark energy). Since we are describing absorption, ($\alpha_* > 0$) this means that the parameters ϵ and η_0 must be positive.

3. Extended Luminosity Distance and Supernova Dimming

Nowadays, constraints based on SNe Ia data are considered the best method for studying the cosmic expansion history at $z < 1.5$. Let us now confront the extended luminosity distance including absorption as an optical metric phenomenon with the latest Supernova data. In our subsequent analyzes we consider only the flat Λ CDM model for which the dimensionless Hubble parameter, $E(z)$, takes the following form:

$$E(z) = \sqrt{\Omega_M(1+z)^3 + \Omega_\Lambda}, \quad (9)$$

where $\Omega_\Lambda = 1 - \Omega_M$.

To begin with, let us consider the Union2 supernova sample which is formed by 557 measurements of distance moduli from SNe Ia as compiled by Amanullah *et al.* (2010). In order to avoid effects from Hubble bubble, only 506 supernovae with redshifts greater than $cz = 7000 \text{ km/s}$ were selected (Conley *et al.* 2007, Kessler *et al.* 2009, Sinclair *et al.* 2010). As widely known, the SNIa Union2 data are obtained by adding new datapoints (including the high redshift SNIa) to the original Union data (Kowalski *et al.* 2008). For this enlarged sample, a number of refinements to the original Union analysis chain has been done, in particular, the relative importance of systematic effects was highlighted (in this connection see also Sullivan *et al.* (2011) to the Legacy Survey sample (SNLS3)).

In our statistical analysis we consider a maximum likelihood determined by a χ^2 statistics

$$\chi^2(\mathbf{p}) = \sum_{SNIa} \frac{[\mu_{obs}(z_i; \mathbf{p}) - \mu_{th}(z_i)]^2}{\sigma_{obs,i}^2}, \quad (10)$$

where $\mu_{th}(z_i) = 5 \log_{10} D_L(z_i, \mathbf{p}) + \mu_0$ is the theoretical distance moduli, $\mu_0 = 25 - \log_{10} H_0$, D_L is the luminosity distance from Eq. (6), $\sigma_{obs,i}$ is the uncertainty in the individual distances (including systematic errors), and the complete set of parameters is given by $\mathbf{p} \equiv (H_0, \Omega_M, \alpha_*)$. It should be stressed that for the considered SNe Ia subsample we have combined the statistical plus systematic errors in quadrature as compiled in Table 7 of Amanullah *et al.* (2010) and neglected σ_{lc}^2 . We have also marginalized on the Hubble constant.

In Fig. 1(a) we show the contours on the $\Omega_M - \alpha_*$ plane corresponding to a flat Λ CDM

model and by considering that the absorption α_* is constant. As indicated, the shadow lines are cuts in the regions of 68.3%, 95.4% and 99.7% of probability. The constraints on the free parameters are restricted to $0.0 \leq \alpha_* \leq 1.55$ and $0.21 \leq \Omega_M \leq 1.0$ at 2σ of statistical confidence while to the latter we have obtained $0.0 \leq \alpha_* \leq 1.46$ and $0.22 \leq \Omega_M \leq 1.0$ at 2σ of statistical confidence. In the absence of absorption ($\alpha_* = 0$) the limit on the density parameter is $0.21 \leq \Omega_M \leq 0.34$ (concordance model) while for $\Omega_M = 1$ (Einstein de Sitter), we find that the absorption parameter lies on the interval $1.20 \leq \alpha_* \leq 1.54$ (2σ). Note that the positiveness of α_* (absorption), implies that a pure de Sitter model ($\Omega_\Lambda = 1, \Omega_M = 0$) is not allowed by such data. The best fit to the free parameters are $\Omega_M = 0.27, \alpha_* = 0.0$ with $\chi_{min}^2 = 330.5$.

In Fig. 1(b) we display the likelihood distribution functions, $e^{-\chi^2/2}$ of α_* , for the Einstein-de Sitter model ($\Omega_M = 1$) with constant absorption. The blue curve includes statistical + systematic errors, but, for comparison, we have also shown the red curve with only statistical errors. The upper and lower horizontal lines correspond to 68.3%, and 95.4% c.l., respectively. By marginalizing on the Hubble parameter we obtain $\alpha_* = 1.38 \pm 0.08(0.15)$ with 1σ (2σ) of probability and $\chi_{min}^2 = 331.3$. In this model the Universe is always decelerating since $q(z) = q_0 = 1/2$. Instead of dark energy we have a cosmic medium whose absorbing properties is quantified by the dimensionless parameter $\alpha_* \simeq 1.4$ which is responsible for the SNe Ia dimming. The dimensional absorption parameter, $\alpha = \alpha_* H_0 \sim 10^{-4} Mpc^{-1}$, is nearly the same one previously obtained by Chen and Kantowski (2009a).

Let us now consider the deformation function, $\eta(z) = (1+z)^\epsilon$, as adopted by Avgoustidis *et al.* (2009). As shown in the previous section, the associated cosmic absorption parameter in this case is $\alpha_*(z) = 2\epsilon E(z)$. In order to simplify the notation, in what follows we consider the parameter, $\alpha_0 = 2\epsilon$.

In Figure 2(a) we display the corresponding plots for the (Ω_M, α_0) plane to the case of variable absorption, $\alpha_* = \alpha_0 E(z)$. As in Fig. 1(a) the analysis includes statistical plus systematic errors. The confidence region (2σ) in this plane is defined by $0.0 \leq \alpha_0 \leq 0.92$ and $0.20 \leq \Omega_M \leq 1.0$. Again, the best fit is the Λ CDM with best fit $\Omega_M = 0.27, \alpha_0 = 0.0$ with $\chi_{min}^2 = 330.5$.

In Fig. 2(b) we display the likelihood distribution functions for α_0 . Blue and red curves correspond to the same analysis of Fig. 1(b), that is, with and without systematics. To the latter case, we obtain that $\alpha_0 = 0.88 \pm 0.05(0.10)$ with 1σ (2σ) of confidence level. Once again we have $q(z) = q_0 = 1/2$ in the presence of the absorption, but with $\chi_{min}^2 = 335.9$.

It should be stressed that the above results for the $\Omega_M - \alpha_*$ and $\Omega_M - \alpha_0$ planes, based on a subsample of the Union2 SNe Ia data, are strongly suggesting that one of the following

possibilities must be true:

(i) The Universe has an extra dark energy component (Λ) which provides the fuel for the observed accelerating stage ($q_0 < 0, \alpha_* = 0$).

(ii) We live in a decelerating Einstein-de Sitter Universe ($\Omega_M = 1$) endowed with a cosmic absorption mechanism which is responsible for the dimming of the distant Supernovae Ia ($q_0 = 1/2, \alpha_* \neq 0$).

(iii) We are not living in the extreme cases above mentioned, that is, both phenomena (cosmic opacity and Λ) are partially operating in the observed Universe.

In conclusion, by adopting the Gordon-Chen-Kantowski description we have shown how to obtain analytical expressions for any deformed distance duality relation (parametrized by $\eta(z)$ function) in terms of the cosmic absorption parameter and vice-versa. Four specific examples of $\eta(z)$ functions, recently proposed in the literature, were explicitly considered. The associated absorption parameter for each one was explicitly determined and some of its physical implications also were discussed in detail. Two different scenarios with cosmic absorption were proposed and their free parameters constrained by using the SNe Ia data (Amanullah et al. 2010). Working in the framework of a flat Λ CDM model we have found that both scenarios ($\alpha_* = \alpha_0 E(z)$ and $\alpha_* = \text{constant}$) are also compatible with a decelerating Einstein-de Sitter Universe (see Figures 1a and 2a). Finally, as remarked before, in the present study we have neglected any effect due to a possible light refraction phenomenon as described in the original Gordon's description. The general case it will be discussed in a forthcoming communication.

JASL and VTZ are partially supported by CNPq and FAPESP (Brazilian Research Agencies). JVC is supported by CAPES.

REFERENCES

- Amanullah, R., et al. 2010, ApJ, 716, 712
- Avgoustidis, A., Verde, L. & Jimenez, R. 2009, JCAP, 0906, 012
- Avgoustidis, A., Burrage, C., Redondo, J., Verde, L., & Jimenez, R. 2010, JCAP, 1010, 024
- Basset, B. A., Kunz M. 2004, Phys. Rev. D, 69, 101305 [astro-ph/0312443v2]
- Bonamente, M., et al. 2006, ApJ, 647, 25

- Chen B. and Kantowski R. 2009a, Phys. Rev. D, 79, 104007
- Chen B. and Kantowski R. 2009b, Phys. Rev. D, 80, 044019
- Conley A. *et al.* 2007, ApJ, 664, L13
- Daly, R. A., & Djorgovski, S. G. 2003, ApJ, 597, 9
- De Bernardis, F., Giusarma, E., & Melchiorri, A. 2006, IJMPD, 15, 759 [gr-qc/0606029v1]
- De Filippis, E., Sereno, M., Bautz, M.W., & Longo, G. 2005, ApJ, 625, 108
- Etherington, I. M. H. 1933, Phil. Mag., 15, 761; reprinted in 2007, Gen. Relativ. Grav., 39, 1055
- Gonçalves, R. S., Holanda, R. F. L., & Alcaniz, J. S. 2011, arXiv:1109.2790v1 [astro-ph.CO]
- Gordon, W. 1923, Ann. Phys. (Leipzig), 72, 421
- Gurvitz, L. I., Kellerman, K. I., & Frey, S. 1999, A&A, 342, 378
- Hicken, M. *et al.*, 2009, ApJ, 700, 1097
- Holanda, R. F. L., Lima, J. A. S., & Ribeiro, M. B. 2010, ApJL, 722, L233 [arXiv:1005.4458]
- Holanda, R. F. L., Lima, J. A. S., & Ribeiro, M. B. 2011a, A&A Letters, 528, L14 [arXiv:1003.5906]
- Holanda, R. F. L., Lima, J. A. S., & Ribeiro, M. B. 2011b [arXiv:1104.3753]
- Kessler R. *et al.* 2009, ApJS, 185, 32
- Khedekar, S. & Chakaborti, S. 2011, Phys. Rev. Lett. 106, 221301
- Kowalski, M., *et al.* 2008, ApJ, 686, 749
- Li, Z., Wu, P., & Yu, H. 2011, ApJL, 729, L14
- Lima, J. A. S., & Alcaniz, J. S. 2000, A&A, 357, 393 [astro-ph/0003189]
- Lima, J. A. S., & Alcaniz, J. S. 2002, ApJ, 566, 15 [astro-ph/0109047]
- Meng, X-L., Zhang, T-J., & Zhan, H. 2011 [arXiv:1104.2833v1]
- Nair, R., Jhingan, S., & Jain, D. 2011, JCAP, 1105, 023 [arXiv:1102.1065]
- Santos, R. C., & Lima, J. A. S. 2008, Phys. Rev. D 77, 083505 [arXiv:0803.1865]

Sinclair B., Davis T. M. & Haugblle T. 2010, ApJ, 718, 1445

Sullivan, M., et al. 2011, Astrophys. J., 737, 102 [arXiv:1104.1444]v2

Stern, D., Jimenez, R., Verde, L., Kamionkowski, M., & Stanford, S. A. 2010, JCAP, 1002, 008 [arXiv:0907.3149].




Fig. 1.— **a)** Contours in the $\Omega_M - \alpha_*$ plane for 506 supernovae data by considering a flat Λ CDM model with constant absorption parameter, α_* . Note that the Einstein-de Sitter model is now allowed by such data. **b)** The likelihood function of the constant absorption α_* for the Einstein-de Sitter model ($\Omega_M = 1$). The red line takes into account only statistical while the blue curve includes both errors (statistical + systematic). The upper and lower horizontal lines correspond to 68.3% (1σ), and 95.4% (2σ) c.l., respectively. The free parameter is constrained to $\alpha_* = 1.38 \pm 0.08$ (statistical + systematic, 1σ c.l.).




Fig. 2.— **a)** Contours in the $\Omega_M - \alpha_0$ plane for the same 506 supernovae data by considering a flat Λ CDM model with an epoch-dependent absorption, $\alpha_* = \alpha_0 E(z)$, as fixed by the Avgoustidis et al. (2009) deformation function (see sections I and II). The best fit to the pair of free parameters is given by $(\Omega_M, \alpha_0) = (0.27, 0.0)$. **b)** The likelihood functions of α_0 in the case of flat Einstein-de Sitter model. The free parameter is constrained to $\alpha_0 = 0.88 \pm 0.05$ (statistical + systematic, 1σ c.l.).